

# Raldon

Oral solution for drinking water for calves, swine, broilers,  
laying hens turkey rabbits (colistine sulphate)

dox-a

Pharma

**Composition per 1lit:** Active substance  
Colistine sulphate 120.00 g

**Pharmaceutical form:** oral solution for drinking water

**Packaging:** 1l bottle

**Target species:** calves, swine, broilers, laying hens, turkeys, rabbits.

**Indication:** infections caused by germs sensitive to colistine sulphate, in particular bacteria enteritis sustained by Gram-negative: E.Coli, Klebsiella spp., Pseudomonas spp., Salmonella spp., Shigella spp.

**Contra-indication:** do not use in case of known resistance to polymyxines, do not administer to animals with accertained hypersensitive to colistine or to one of the excipients.

**Adverse Reaction:** in rare cases, oral administration, with long and/or excessive doses, may cause gastrointestinal disease (vomit, nausea, diarrrea). If heavy adverse reactions happen or other reaction not mentioned in this label, please inform the veterinary.

**Special warning for each target species:** the assumption of medicated water by animals could be altered due to the disease. For subjects which present a reduces water assumption do the treatment by parental way using correct injectable product upon veterinary suggestion.

**Special precaution in use**

**Special precaution for use in animal:** due to possible variation during time and sensitive geography of the microbial target species to colistine we suggest to do the bacteriological exam and sensitivity test.

Choose antibiotics with action spectra reduced against the one of colistine when antiobiogram confirm the possible efficacy of the treatment. Not correct use of the product may increase the prevalence of germs resistant to colistine and reduce the efficacy of treatment with other antibiotics of the polimixine group caused by possible cross resistance. Do not leave medicated water available to other animals. The use of the veterinary medicine (colistine) in broiler has to be conform to CE1177/2006 regulation of the commission and to national law.

**Special precaution to be adopted by the person who administer the product to animals.** Adopting the usual proceeding for preparation and administration of product particular precaution for operator aren't necessary, however it's better avoid inhalation and direct contact; we suggest use protective gloves for handling product; do not eat, drink and smoke during handling; wash hands after use. In case of contamination well wash with water and soap. If irritation persists apply to the doctor. The persons with well known sensitivity to colistine has to avoid contacts with the veterinary drug.

**Use during pregnancy, lactation and ovodeposition:** particular dangerous effects are not mentioned for this target specie. In any case product has to be administered only in case of effective needs and under veterinary control.

**Interaction with other veterinary medicines and other**

**interaction form:** present cross resistance with Polimixine B.

**Overdosage (symtom emergency actions, antidotes) if necessary**

No toxic effects are known by overdosage related to the use of product. Do not increase suggested dosages.

**Incompatibility.** In absence of compatibility studies this medicine has not to be mixed with other veterinary medicines.

**Dose and administration, treatment time.**

The medicinal product must be administered diluted in drinking water, for 3-15 days as per veterinary prescription, at daily dose 5-6mg colistine sulphate/kg l.w. (41.67 ml- 50ml of product/ 1000kg b. wt./3-15 days).

The correspondant concentration of drinking water has to be recalculated in function of the real animal weight and effective water consumption. Do not over authorised dose in mg of active principle/kg l.w. In order to avoid under dosage the body weight has to be determinate in the most accurately way.

The consumption of medicated water depends from clinical animal conditions.

In order to obtain a correct dosage, concentration of antimicrobial has to be recalculated conformity. Do not mix with solid feed. For a correct administration remove liquid feed residue and administer the medicated one with "colistine sulphate 12% liquid" in the above mentioned quantity. At the end of therapy give again the feed not medicated in order to grand a correct dose administered.

Medicated water consumption depend from clinical conditions of animals.

Use machine for dosage correctly calibrated.

**Withdrawal time:** Meat and offals:

Rabbits 5 days

Swine 7 days

Calves 7 days

Broilers 1 days

Turkeys 1 days

Laying hens: eggs: 5 days

**STORAGE:** store at temperature not exceeding 25°C and after opening used for 30 days at the same temperature and after dilution used for 24 hours.

**Special precautions for the disposal of unused products or waste materials**

The unused product or waste from those medicated products must be delivered to the usual collection points for expired or unused medicines in conformity to disposition given by local law.

**Keep away from children reach.**

**FOR VETERINARY USE ONLY**

Net weight 1L

REGISTRATION N. 4033/2017

Batch No.: SEE ON THE CAP

Mfg date: SEE ON THE CAP

Exp. date: SEE ON THE CAP

**ELYOSER**  
Medicine Trading  
اليسر تميز العصر